

Comparative anatomy of carpal and tarsal bones in three sympatric Turolian hipparions (Kryopigi, Chalkidiki, Greece)

Georgios LAZARIDIS & Evangelia TSOUKALA

G. Lazaridis, Aristotle University, Geology School, GR-54124 Thessaloniki, Greece; geolaz@math.auth.gr
E. Tsoukala, Aristotle University, Geology School, GR-54124 Thessaloniki, Greece; lilits@geo.auth.gr

ONLINE SUPPLEMENTARY INFORMATION

Various hipparion groups of the Kryopigi locality are separated in the scatter plots of Figs 1 and 2. The separation is based on their dimensions and specimens found articulated with metapodials and other long bones are used as a reference.




Fig. 1 - Scatter-diagrams of the Kryopigi hipparion carpal bones. Open square: *C. matthewi/nikosi*; solid square: *H. phlegrae*; triangle: *?Cremohipparion* sp.; asterisk: *H. cf. H. brachypus*. H: height; DAP: diameter anteroposterior; DT: diameter transverse.




Fig. 2 - Scatter-diagrams of the Krypigi hipparion tarsal bones. Square: *C. matthewi/nikosi*; solid square: *H. phlegrae*; triangle: *?Cremohipparion* sp.; asterisk: *H. cf. H. brachypus*. H: height; DAP: diameter anteroposterior; DT: diameter transverse.