

Recent infralittoral Foraminiferida and Ostracoda from the Porto Cesareo Lagoon (Ionian Sea, Mediterranean)

Giuseppe AIELLO, Diana BARRA, Maria Grazia COPPA, Assunta VALENTE & Francesco ZENI

- G. Aiello, Dipartimento di Scienze della Terra, Università degli Studi di Napoli "Federico II", Largo S. Marcellino 10, I-80138 Napoli, Italy; aie64llo@hotmail.com
D. Barra, Dipartimento di Scienze della Terra, Università degli Studi di Napoli "Federico II", Largo S. Marcellino 10, I-80138 Napoli, Italy; dibarra@unina.it
M.G. Coppa, Dipartimento di Scienze della Terra, Università degli Studi di Napoli "Federico II", Largo S. Marcellino 10, I-80138 Napoli, Italy; grappa@unina.it
A. Valente, Dipartimento di Scienze della Terra, Università degli Studi di Napoli "Federico II", Largo S. Marcellino 10, I-80138 Napoli, Italy; valente.assunta@libero.it
F. Zeni, Dipartimento di Scienze della Terra, Università degli Studi di Napoli "Federico II", Largo S. Marcellino 10, I-80138 Napoli, Italy; francesco.zeni@tin.it

KEY WORDS - *Benthic Foraminiferida, Ostracoda, Recent, Gulf of Taranto, Infralittoral assemblages.*

ABSTRACT - *Recent benthic Foraminiferida (111 species, 45 genera) and Ostracoda (150 species, 43 genera) assemblages were studied. Samples were collected in the southern part of the Porto Cesareo Lagoon, on the south-western coast of Apulia, Ionian Sea. Thirty-two bottom samples were collected by hand at a water depth of 0.12 to 4.78 m. Assemblages are characteristic of marine infralittoral warm waters, in agreement with the previous records of benthic organisms with tropical affinity. Dominant taxa are either phytophilous or typical of sandy bottom. Abundance and diversity are controlled by environmental factors such as the type of substrate, hydrodynamic conditions and distance from the open sea. Anthropic pollution negatively influences microfaunal assemblages only for a few dozens of meters away from the urbanized area.*

RIASSUNTO - [Foraminiferi ed ostracodi di ambiente infralittorale della Laguna di Porto Cesareo (Mar Ionio, Mediterraneo)] - *Sono state studiate le associazioni a foraminiferi bentonici e ad ostracodi provenienti da 32 campioni di fondo raccolti manualmente tra 0,12 e 4,78 metri di profondità nell'insenatura della Strea, parte della Laguna di Porto Cesareo, situata sulle coste ioniche della Puglia. Le faune sono composte da 111 specie di foraminiferi bentonici appartenenti a 45 generi e 150 specie di ostracodi, ripartite in 43 generi. Le associazioni sono caratteristiche di acque marine infralittorali temperato-calde, in accordo con la già nota presenza di organismi ad affinità tropicale e rappresentate da specie tipiche di substrati sabbiosi e specie fitofile. La distribuzione delle microfaune si è dimostrata sensibile principalmente a fattori quali il tipo di substrato, l'idrodinamismo e la distanza dall'imboccatura della laguna. Inoltre gli apporti antropici inquinanti esercitano un'evidente influenza sui microorganismi ma limitatamente alle acque immediatamente prospicienti l'area più intensamente urbanizzata.*

INTRODUCTION

This study was undertaken to define the foraminiferal and ostracod assemblages in a sheltered inlet, connected to the Ionian Sea (Fig. 1). The environment of this small bay, part of the Porto Cesareo Lagoon, is characterized by very shallow, well oxygenated, and low polluted marine waters. The calm water conditions allowed the good preservation of the thanatocoenosis, including delicate shells which are generally not preserved in high energy upper infralittoral environments.

The study area is located on the Ionian coast of the Salentine peninsula (Apulia), about 60 km east of Taranto town and 24 km south-west of Lecce (lat. 40° 15' N, long. 17° 54' E). The La Strea Bay ("Laguna di Porto Cesareo" in Passeri, 1974; "Insenatura della Strea" in Belmonte & Rubino, 1988; "Baia della Strea" in Castelli et al., 1988) is an approximately elliptical

inlet 2500 m long with a maximum width of about 1000 m. It has been formed behind an ancient barrier island system which is currently represented by the peninsula of La Strea, which bounds the bay on the south-west, and some islets on the north-west. It was connected to the Ionian Sea in the last phase of the Holocene transgression, which turned the ancient marsh into an euhaline shallow inlet (Passeri, 1974; Mastronuzzi et al., 1989). The channel system, which connect the bay with the open sea, is situated in the north-western part. It is about 700 metres wide. Supratidal marshes, described by Passeri (1974), are now reclaimed.

The bottom consists mainly of unconsolidated coarse- to fine-grained sands, with a carbonate content ranging from 53 to 90% (Passeri, 1974; Ricchetti & Nuovo, 1988).

The waters are very shallow, the average depth being about one meter. Maximum water depth, in the north-western portion of the bay, is about 4.8 m. The




Fig. 1 - Location map of the study area, showing transects and sampling stations.

maximum depth of the channels connecting the bay to the sea is 2.5 m (Passeri, 1974). Tidal range is about 20–30 cm.

The water temperature varies from a low of 9.2°C in February to a high of 28.2°C in August (Belmonte & Rubino, 1988; Mercurio et al., 2000), showing a sharp increase during summer (Gherardi et al., 2001).

The salinity of the La Strea Bay is slightly lower (about 1‰) than the open sea waters, due to karstic freshwater supplies along the coast. Nonetheless, the bay water is euhaline (33.5 to 37.8‰). Dissolved oxygen content and pH range from 6.2 to 10.9 ppm and 7.7 to 8.7, respectively.

The waters in the vicinity of the township of Porto Cesareo show a relatively high concentration of nitrites and nitrates, indicating organic pollution due to human activities. The annual mean values of nitrites and nitrates measured by Belmonte & Rubino (1988) are noticeably higher in the northernmost stations (nitrites, 0.567 µg-atN/l; nitrates 19.79 µg-atN/l) than in both the central part of the bay (nitrites = 0.251 µg-atN/l, nitrates = 31.99 µg-atN/l) and in open sea (nitrites = 0.193 µg-atN/l, nitrates = 15.61 µg-atN/l).

The aquatic vegetation is dominantly a mixed meadow of *Cymodocea-Caulerpa*. On the northern edge of the bay, i.e. close to the town, nitrophilic taxa such as *Ulva* and *Enteromorpha* are present. Thermophilic, photophilous assemblages, including the chlorophyte *Anadyomene stellata*, a species with tropical affinity, live in the very shallow, calm waters of the southwestern part of the bay, on rocky bottoms (Pardi et al., 1988).

Parenzan (1983) emphasized the meaning of the subtropical biocoenosis defined by the co-occurrence of *A. stellata*, the demosponge *Geodia cydonium* and

the echinoderm *Holothuria impatiens*. Noteworthy is the presence of the warm water sabellid *Fabricia filamentosa* and the gastropod *Pirenella conica* (Castelli et al., 1988).

MATERIALS AND METHODS

A total of 32 sediment samples was collected in the La Strea Bay. Twenty-eight along six parallel transects plus four “isolated” samples. The maximum water depth from which samples were collected was 4.78 m; the minimum was 0.12 m. Core samples were obtained by direct insertion of a short core tubes (h = 12 cm; Ø = 8 cm) into the sediment. After core recovery, subsamples of the surface sediments were collected from the central (undisturbed) part of core, to analyze recent (subrecent) assemblages. Bathymetry, grain size, number, location, and dimension of samples are reported in Fig. 1 and Tab. 1.

The samples were wet-sieved through 170-mesh (90 µm) screens and oven-dried. Residue granulometry was determined by using a stereo microscope fitted with an ocular micrometer. Sediment classification was based on Wentworth grain-size scale.

Biogenic components consist mainly of foraminifers and subordinately of bryozoans fragments, sponge spicules, bivalves and gastropods (both shells and fragments), ostracods, echinoid spines, annelids.

A microsplitter was used to obtain a split of approximately 300 specimens; due to the paucity of

Samples	Longitude E	Latitude N	Depth (m)	Thickness (cm)	Dry weight (gr)	Residue weight (> 90 µm)	Grain size
A1	17°53.67'	40°15.92'	0.43	4	30.9	27.9	FS
A2	17°53.76'	40°15.83'	1.30	3.5	23	16.7	MS-FS
B1	17°53.57'	40°15.64'	3.21	3	40.1	19	CS-FS
B2	17°53.64'	40°15.71'	2.43	4	39.9	31.2	CS-MS
B3	17°53.73'	40°15.78'	2.40	4	36.1	22.8	CS
B4	17°53.87'	40°15.88'	1.40	3	24.1	21.5	CS
B5	17°53.93'	40°15.91'	0.90	3	31.2	23.1	CS-MS
C1	17°53.52'	40°15.34'	4.40	2.5	34	27.2	CS
C2	17°53.61'	40°15.40'	4.78	3.5	23.9	20.5	CS
C3	17°53.72'	40°15.46'	4.50	5	32	21.6	CS-FS
C4	17°53.83'	40°15.53'	3.40	3	26	18.8	CS
C5	17°53.92'	40°15.61'	2.58	3.5	33	23.2	CS-FS
C6	17°54.02'	40°15.66'	1.73	4.5	29	24.2	CS
C7	17°54.10'	40°15.72'	-	3.5	42.1	34	CS
D1	17°53.85'	40°15.33'	1.52	4	32	22.5	MS-FS
D2	17°54.01'	40°15.38'	2.30	7	36.2	25.9	CS-MS
D3	17°54.17'	40°15.49'	1.51	4	25.9	18.2	CS-FS
D4	17°54.32'	40°15.59'	1.11	3	26.9	18.5	CS-FS
E3	17°54.01'	40°15.07'	1.42	3.5	35.2	19.8	MS-FS
E4	17°54.25'	40°15.16'	1.50	4	31.8	21	CS-MS
E5	17°54.26'	40°15.23'	1.49	3	34.1	21.2	FS
E6	17°54.45'	40°15.28'	1.30	3.5	34.2	23.8	MS-FS
F1	17°54.28'	40°14.84'	0.50	2.5	36.8	25.1	CS-FS
F2	17°54.34'	40°14.88'	0.94	4	30	23.1	MS-FS
F3	17°54.45'	40°14.97'	1.12	5.5	32.5	14	FS
F4	17°54.60'	40°15.04'	1.50	3	30.9	22	MS-FS
G1	17°54.47'	40°15.63'	0.20	2.5	33	24	MS-VFS
G2	17°54.52'	40°14.67'	0.70	3	38.4	29	MS
G3	17°54.57'	40°14.71'	0.80	2.7	31.7	22	MS-VFS
G4	17°54.66'	40°14.77'	0.65	3.5	35.1	26.9	MS-VFS
H	17°54.69'	40°14.61'	0.26	3.5	31	22.8	VFS
I	17°54.69'	40°14.51'	0.12	4.5	30.2	25.7	FS

Tab. 1 - Coordinates, depth, thickness, weight and granulometry of the studied samples.




Fig. 2 - Foraminifer abundance (A = number of specimens per gram of dry sediment) and diversity (SR = number of species, SI = Shannon index). Horizontal line indicates medium value.

ostracod shells, a total picking was carried out for 16 samples. The species were identified and the adult specimens counted.

RESULTS AND DISCUSSION

Foraminiferida (M.G. Coppa, A. Valente & F. Zeni)

One-hundred twelve foraminiferal species, 111 benthic and one planktic, have been recorded (Tab. 2). The systematics follows the scheme proposed by Loeblich & Tappan (1964, 1987).

The assemblages consist mainly of well-preserved benthic foraminifers: *Peneroplis pertusus* (26.87-61.10%) and, subordinately, *Siphonaperta agglutinans* (0.23-16.37%), *Peneroplis planatus* (0.26-12.37%), *Ammonia beccarii* (0.49-11.11%), *Sardamina cherchiai* (0.29-10.27%), *Elphidium macellum* (0.28-9.09%), *E. advena* (0.29-9.31%), *Spirolina acicularis* (0.25-8.09%), *Quinqueloculina parvula* (0.33-4.60%), *Q. seminulum* (0.26-3.83%). In the southern, sheltered part of the lagoon, beside the above mentioned species *Haynesina depressula* (0.20-5.77%), *Glabratella patelliformis* (0.21-5.42%), *Ammonia tepida* (0.24-3.24%) also occur. Rare planktic foraminifer specimens, all referable to *Neogloboquadrina pachyderma*, are poorly preserved.

Near the opening of the lagoon to the sea, on coarse sediments, in deeper, high energy waters, benthic foraminifers are mainly large, thick and abraded. In the northern and southern sheltered areas, on fine sediment substrate, in very shallow, low energy waters, foraminifers are smaller, delicate and generally well-preserved. Part of the assemblages, ranging from 1.97

to 25.56%, are indeterminable due to the poor state of preservation.

Foraminifer abundance variability is higher than the diversity variability (Fig. 2). Abundance ranges from 72 (sample C1) to 9297 (sample E6). The lowest diversity is recorded in the samples C1 (SR = 22) and B2 (SI = 1.78), the highest in the samples B1 (SR = 60) and F3 (SI = 3.24).

ASSEMBLAGE INTERPRETATION

Present data have been compared with the ones previously reported for the Mediterranean waters by Le Calvez & Le Calvez (1958), Blanc-Vernet (1969, 1974, 1988), Blanc-Vernet et al. (1979), Cita & Premoli-Silva (1967), Sgarrella & Barra (1985), Sgarrella et al. (1985), Coppa (1987, 1991), Langer (1988), Scorziello (1990), Vè nec-Peyrè & Le Calvez (1988), Murray (1991), Sgarrella & Moncharmont Zei (1993), Coppa et al. (1994), and Coppa & Di Tuoro (1995), with the aim to confirm the ecological meaning of the assemblages and to indicate new distribution data.

Taxa whose ecological characteristics are known could be ascribed to the groups epiphyte, euryhaline, sandy bottom dwellers and warm-temperate species (Fig. 3).

- Epiphyte species: *Nubecularia lucifuga*, *Peneroplis pertusus*, *P. planatus*, *Spirolina acicularis*, *Discorbis mira*, *Neoconorbina terquemi*, *Rosalina bradyi*, *R. globularis*, *R. floridana*, *Glabratella patelliformis*,




Fig. 3 - Percent values of foraminifer ecological groups.

[illegible]

Tab. 2 - Percent values of foraminifer species.

SPECIES		/		SAMPLES																		
				depth (m)																		
Articulina carinata Wiesner																						
Wellmanirella striata (Sidedbottom)																						
SORITIDAE																						
Sporollina aculeatus (Bartsch)																						
Penorthis peratus (Forsk.)																						
Peneroplis planatus (Fichtel & Moll)																						
NODOSARIIDAE																						
Oreomorphina sp.																						
POLYMORPHINIDAE																						
Polymorphina gibba (d'Orbigny)																						
Polymorphina sp.																						
BOLIVINITIDAE																						
Bolivina purcidata d'Origny																						
Bolivina spatulata (Williamson)																						
BULIMINIDAE																						
Bulinirella basiscostata Parr																						
Bulinina pupoides d'Orbigny																						
Bulinina sp.																						
UWIGERINIDAE																						
Uvigerina mediterranea Holker																						
Uvigerina sp.																						
DISCORBIDAE																						
Buccella granulata (Di Napoli)																						
Buccella sp.																						
Discorbirella bergheloti (d'Orbigny)																						
Discorbis mira Cushman																						
Discorbis aff. mira Cushman																						
Neoncorbina terquemi (Rzehak)																						
Neoncorbina posidonicola (Colom)																						
Rosalina bradyi Cushman																						
Rosalina concinna (Brady)																						
Rosalina floridana (Cushman)																						
Rosalina globularis d'Orbigny																						
Rosalina obtusa d'Orbigny																						
Rosalina cf. viarebana d'Orbigny																						
Rosalina sp.																						
Valvulineria minuta Parker																						
GLABRATELLIDAE																						
Glabratella erecta (Sidedbottom)																						
Glabratella hexacamera (Seiglie & Bermudez)																						
Glabratella imperatoria (d'Orbigny)																						
Glabratella patelliformis (Brady)																						
Glabratella torrei (Bermudez)																						
ASTERIGERINIDAE																						
Astigeriginata cf. maritima (Williamson)																						
ROTALIDAE																						
Ammonia beccarii (Linnee)																						
Ammonia gaimardi (d'Orbigny)																						
Ammonia tepida Cushman																						
Ammonia sp.																						
ELPHIDIIDAE																						
Elphidium aculeatum (d'Orbigny)																						
Elphidium advenum (Cushman)																						
Elphidium crispum (Linnee)																						
Elphidium cuvillieri (Levy)																						
Elphidium macellum (Fichtel & Moll)																						
Elphidium sp.																						
EPONIDAE																						
Eponides sp.																						
CIBICIDIDAE																						
Cibicides variabilis (d'Orbigny)																						
Cibicides lobatulus (Walker & Jacob)																						
Cibicides relictus Montfort																						
Cibicides sp																						
PLANORBULINIDAE																						
Planorbulina mediterraneensis d'Orbigny																						
CYMBALOPORIDAE																						
Cymbaloporella bradyi (Cushman)																						
NONIONIDAE																						
Nonion depressulum (Walker & Jacob)																						
Nonion sp.																						
Nonionella optima Cushman																						
GLOBOROTALIDAE																						
Globorotalia inflata (Ehrenberg)																						

Tab. 2 (continue)

Elphidium crispum, *E. macellum*, *Lobatula lobatula*, *Cibicides refulgens*.

- Euryhaline species: *Ammonia beccarii*, *A. tepida*, *Haynesina depressula*.

- Sandy bottom species: *Sardammina cherchiaie*, *Quinqueloculina berthelotiana*, *Q. bosciiana*, *Q. bradyana*, *Q. parvula*, *Q. seminulum*, *Q. stalker*, *Q. stelligera*, *Q. striata*, *Siphonaperta agglutinans*, *S. aspera*, *Pseudotriloculina oblonga*, *Triloculina labiosa*, *Valvulineria minuta*, *Ammonia beccarii*, *A. gaimardi*, *A. tepida*, *Elphidium advena*, *E. crispum*, *E. macellum*, *Haynesina depressula*.

- Warm-temperate species: *Peneroplis pertusus*, *P. planatus*, *Spirolina acicularis*, *Glabratella patelliformis*.

Overall, the foraminiferal samples of the Porto Cesareo lagoon are characterized, mainly in the southern (samples F4 and G1) and northern (samples B4, C1, C2) part of the bay, by the dominance of some taxa. These are eurythermal and euryhaline taxa (peneropliids, miliolids, nonionids, *Ammonia*) tolerating the environmental variability of the lagoon (Peres & Picard, 1964; Carrada & Fresi, 1988).

In the southern part of the lagoon it has been noted that euryhaline species are relatively more abundant (0.81% - 15.32%). The distribution pattern of these taxa could be due to the freshwater supply mentioned by Passeri (1974).

Abundance in peneropliids and glabratellids (28.95% - 73.48%) confirms their preference for warm waters (Phleger, 1960; Seiglie & Bermúdez, 1965) and lagoonal environments (Cita & Premoli-Silva, 1967; Murray, 1991). The abundance of *Peneroplis pertusus*, a species commonly living on firm sandy bottoms and on seagrasses, in shallow, warm waters (Blanc-Vernet, 1969; Colom, 1971; Scorziello, 1990; Murray, 1991) confirms the above mentioned subtropical affinity.

The common occurrence of epiphyte species, ranging from 43.50 to 79.82%, reflects the presence of rich and well diversified vegetation (Pardi et al., 1988).

Sandy bottom species show percent values from 0.80 to 39.07%, generally less abundant than epiphyte species.

The frequency of species typical of deeper waters such as *Sardammina cherchiaie*, *Quinqueloculina stalker*, *Pseudotriloculina oblonga*, *Valvulineria minuta*, suggests an enlargement of their habitat (Coppa, 1991; Sgarrella & Moncharmont-Zei, 1993).

Poorly preserved, abraded specimens, reaching very high values (45.32%) in the sample D4, seem to be indicative of high-energy environment there. It cannot be excluded they are partially transported.

The presence of *Ammonia gaimardi* at the mouth of the lagoon (samples C1, C2), on coarse grained sediments, and near the urbanized area (sample B4), confirms its preference for coarse sandy bottom conditions (Blanc-Vernet, 1969) and its tolerance of organic-rich (Sgarrella & Moncharmont-Zei, 1993) and polluted environments (Sgarrella & Barra, 1985).

Species distribution, shape and dimension of the foraminifers are different between the sheltered and open areas, probably because of different hydrodynamic




Fig. 4 - Percent values of foraminifers characteristic of sheltered and open zone.

regimes. Three groups of species have been defined accordingly (Fig. 4):

- Open zone species (0.46% - 27.48%): *Quinqueloculina berthelotiana*, *Rosalina floridana*, *Ammonia gaimardi*, *Elphidium crispum*.

- Sheltered zone species (0.93% - 27.27%): *Quinqueloculina seminulum*, *Q. striata*, *Siphonaperta aspera*, *Glabratella patelliformis*, *Ammonia tepida*, *Elphidium advena*, *Cibicides refulgens*, *Haynesina depressula*.

- Ubiquitous species (40.93% - 83.24%): *Sardammina cherchiaie*, *Quinqueloculina bradyana*, *Q. parvula*, *Q. stelligera*, *Siphonaperta agglutinans*, *Peneroplis pertusus*, *P. planatus*, *Spirolina acicularis*, *Ammonia beccarii*, *Elphidium macellum*.

Ostracoda (G. Aiello & D. Barra)

The ostracod fauna consists of 150 species, belonging to 43 genera. Forty-two species have been tentatively identified or left in open nomenclature because of sparse, poorly preserved material or due to the absence of adult specimens. Identification of species is based on classical and modern literature on recent ostracods of the Mediterranean (*i.a.* Müller, 1894; Barbeito-Gonzalez, 1971; Uffenorde, 1972; Bonaduce et al., 1976; Breman, 1976; Yassini, 1979; Barra, 1997).

The species list and their quantitative distribution are given in Tab. 3.

In general, the ostracods are well preserved; both adult and young instars shells (carapaces or valves) are present.

The ostracod assemblages are dominated by the genera *Semicytherura* Wagner, 1957 (26 species), *Paradoxostoma* Fischer, 1855 (13 species), *Callistocythere* Ruggieri, 1953 e *Cytherois* G.W. Müller, 1894 (11 species), *Loxoconcha* G.O. Sars, 1866 (10 species), *Neonesidea* Maddocks, 1969 (7 species), *Leptocythere* G.O. Sars, 1925, and *Microcythere* G.W. Müller, 1894 (6 species). The most abundant species are *Xestoleberis dispar* (1130 specimens), *X. communis* (857), *Cistacythereis* (H.) *turbida* (830), *Loxoconcha gibberosa* (626), *Urocythereis distinguenda* (585), and *Loxoconcha stellifera* (482). Further common species (200-400 specimens) are, in decreasing order of

SPECIES / SAMPLES	A1	A2	B1	B2	B3	B4	B5	C1	C2	C3	C4	C5	C6	C7	D1	D2	D3	D4	E3	E4	E5	E6	F1	F2	F3	F4	G1	G2	G3	G4	H	I	
depth (m)	0.43	1.30	3.21	2.43	2.40	1.41	0.90	4.38	4.78	4.46	3.40	2.58	1.73	n.o.	1.52	2.30	1.51	1.11	1.42	1.54	1.49	1.30	0.50	0.94	1.12	1.50	0.20	0.70	0.80	0.65	0.26	0.12	
<i>Agliocypris complanata</i> (Brady & Robertson, 1868)			0.59							1.22							1.34												0.40	0.46			
<i>Agliocypris rara</i> G.W. Müller, 1894						3.33																											
<i>Aurilla arborea</i> (Brady, 1865)	4.00	0.67	1.18	3.54	3.33						0.41	1.17	1.23	1.23	1.81	0.41	2.01	0.44	0.34			0.45			0.85	0.65	0.41	0.40		0.93		1.16	
<i>Aurilla convexa</i> (Brady, 1865)	4.00	6.04	1.18	4.55	7.96	10.00	17.14	7.14	12.50	1.22	2.88	0.58	1.05	10.43	4.07	0.53	0.82	4.03	0.87		0.35	0.45	1.18	1.92	0.43		0.82	1.62	1.21	0.46	1.09		
<i>Aurilla glyptica</i> Barbelto-Gonzalez, 1971					2.86					0.81											0.35		0.59	0.96					1.21		1.45	1.16	
<i>Aurilla prasina</i> Barbelto-Gonzalez, 1971						3.33				1.22	3.29			0.61	0.90	1.06				0.35	0.45	0.59	0.96	0.43			1.23						
<i>Aurilla punctata</i> (Münster, 1830)	4.00	7.38	2.94	22.73	11.50	3.33	12.86	35.71	12.50	0.81	1.65	1.75	5.26	1.23	1.81	2.12	7.76	3.36	1.31	1.35	0.35	0.45	0.59	0.96	0.43		0.41	1.21	0.40	0.46	0.36	1.16	
<i>Basslerites berchioni</i> (Brady, 1870)			0.59	2.27	0.88					0.41	0.41	0.58			1.36	0.53			2.18	2.03	0.71	0.90	2.96	0.96	3.83	1.95		3.64			0.58		
<i>Callistocythere adriatica</i> Masoli, 1968																																	
<i>Callistocythere folliculosa</i> Bonaduce et al., 1976										1.22		1.05																					
<i>Callistocythere aff. C. gliva</i> Bonaduce et al., 1976		2.68	2.35				1.43			0.81	1.23	3.51	3.16	1.84	0.45	3.17	3.67	0.67	1.75	0.68	0.71	0.90							0.40	2.31	3.26	4.65	
<i>Callistocythere intricatoides</i> (Ruggieri, 1953)		1.34	6.47	0.88						1.63																			0.40	0.46	0.72	1.74	
<i>Callistocythere litoralis</i> (G.W. Müller, 1894)																																	
<i>Callistocythere lobanica</i> (G.W. Müller, 1894)			0.59									0.58									0.35												
<i>Callistocythere aff. C. pallida</i> (G.W. Müller, 1894)										0.81	1.23	1.17		1.84						0.35													
<i>Callistocythere aff. C. rastifera</i> (Ruggieri, 1953)		0.67	1.18	0.88		1.43						4.21								0.71								0.41					
<i>Callistocythere sp. A</i>																																	
<i>Callistocythere sp. B</i>			4.55																														
<i>Callistocythere</i> spp.						3.33	1.43				0.82																		0.40				
<i>Cannocythereis whitei</i> (Baird, 1850)										1.23		2.34																	1.62	0.41	1.62	0.58	
<i>Caudites calceolatus</i> (Costa, 1853)	4.00	0.67	0.59		0.88					0.82					2.26	0.53		0.44											0.81		0.36		
<i>Cistacythereis (H.) jurida</i> (G.W. Müller, 1894)	4.00	3.36	1.18	4.55	2.65		2.86	7.14		0.81	5.76	11.70	3.16	1.84	1.36	3.70	7.76	2.68	11.79	13.18	13.83	13.00	8.28	9.62	8.51	18.18	2.88	10.93	11.34	10.65	5.80	8.14	
<i>Costa biter</i> (Brady, 1866)										0.41						0.53														0.46			
<i>Cyprioides torosa</i> (Jones, 1850)		0.67								0.41	0.82	0.58	0.61	0.45	0.53				0.34		0.35		1.18	0.96	0.43	0.65	0.82	0.81	0.40		0.72	1.74	
<i>Cythereoides sordida</i> (G.W. Müller, 1894)			1.18				1.43			0.41	0.41	0.58	1.05	0.61	0.90	2.65	0.82	2.01	2.18	3.04	3.19	2.24	1.18	2.88	1.70	1.95	0.82	2.02	2.43	3.24	1.45	4.65	
<i>Cytherea adriatica</i> Ruggieri, 1952		6.71	0.59	6.82	7.08	3.33	4.29	21.43	25.00	0.41	2.88	7.60	1.05	0.61	2.26	3.17	1.63	0.67	2.18	4.39	1.06	1.79	0.59	2.88	0.43	2.60	0.41	2.02	1.21	2.78	4.35	2.91	
<i>Cytherea subrotunda</i> (Roemer, 1838)	2.68		4.55	1.77	2.86					1.65		2.92			0.45	1.59	2.45	1.34	0.67	2.36	2.13	1.35	0.59	5.77	0.85	4.55	1.23	3.24	2.43	1.39	1.81	0.58	
<i>Cythereis aff. C. fischeri</i> (Sars, 1866)																																	
<i>Cythereis frequens</i> G.W. Müller, 1894												1.05										0.45											
<i>Cythereis aff. C. frequens</i> G.W. Müller, 1894										0.41					0.53				0.34														
<i>Cythereis incongruens</i> G.W. Müller, 1894										0.41		2.34																					
<i>Cythereis aff. C. incongruens</i> G.W. Müller, 1894										0.81		4.68			2.12	0.41	4.70	3.06	1.35		3.19	1.79						0.41					
<i>Cythereis jachinot</i> Barra, 1992		0.67	2.94	0.88											1.06	0.41	0.67	0.44										0.41					
<i>Cythereis pontica</i> Marinov, 1966		0.67	1.18							0.41		0.58																					
<i>Cythereis succinea</i> G.W. Müller, 1894																																	
<i>Cythereis aff. C. triangularis</i> Bonaduce et al., 1979															0.41						0.35												
<i>Cythereis aff. C. offenordii</i> Ruggieri, 1975																													0.81	0.46			
<i>Cythereis</i> sp.										0.41		1.17			0.45	1.06	1.22	1.34	0.44		0.71												
<i>Cythereoma</i> cf. <i>C. karadagiensis</i> Dubowsky, 1939																																	
<i>Cythereoma variabilis</i> G.W. Müller, 1894																																	
<i>Eufonia</i> minima (Bonaduce et al., 1976)		0.59				1.43				1.22	0.41			1.23	0.53			1.34			0.71												
<i>Eucythere curta</i> Ruggieri, 1975																		0.67															
<i>Falsocythere lenyi</i> (Holden, 1967)															0.90																		
<i>Hemicytherea deflori</i> Ruggieri, 1953										0.41																							
<i>Hemicytherea vitensis</i> (G.W. Müller, 1894)			4.71							3.25	0.82	0.58	4.21	1.84	1.36	1.06	0.41	5.37	1.31	0.34	0.35				0.43	0.65	0.82						
<i>Heterocythereis voraginosus</i> Athersuch, 1979										0.41				3.07	0.53																		
<i>Leptocythere lagunae</i> Hartmann, 1958	2.01			0.88		1.43				0.82	0.58	0.61	0.58	0.61	0.53				0.87	0.68			0.59		0.85	0.41	0.40			0.36			
<i>Leptocythere levis</i> (G.W. Müller, 1894)										0.41	0.82	0.58	0.61	0.58	1.59	1.22			2.62	3.38	10.99	5.83	11.24	3.85	4.26	1.95	11.11	6.07	19.03	11.57	1.09	3.49	
<i>Leptocythere ramosa</i> (Rome, 1942)										0.41		0.58																					
<i>Leptocythere aff. L. rara</i> (G.W. Müller, 1894)													2.11																				
<i>Leptocythere</i> sp. 1																																	
<i>Leptocythere</i> sp.																																	
<i>Leptocythere</i> "sp."			2.27																														
<i>Loxocantha affinis</i> (Brady, 1866)		1.34	5.88		0.88					1.65				7.98	5.43	8.47	2.04		2.62	2.03	3.55		4.14		1.28		5.76	2.83	4.63	5.43			
<i>Loxocantha gibberosa</i> Terquem, 1878	4.00	4.03	8.82	6.82	0.88	10.00	2.86			6.50	9.88	2.34	2.11	4.91	7.24	4.23	7.35	2.68	8.73	6.76	7.80	4.93	4.73	0.96	23.40	5.84	1.65	3.64	1.62	2.78		5.81	
<i>Loxocantha aff. L. gibberosa</i> Terquem, 1878										1.23																							
<i>Loxocantha ovulata</i> (Costa, 1853)	4.03		0.59	5.31		1.43				0.81	4.12	9.36		0.61	2.26	2.65	1.63		0.44	0.34	0.90			0.96			0.40				0.58		
<i>Loxocantha pontica</i> Kile, 1937										0.82	0.58	0.58	3.07	3.07	1.06	1.22			0.44	0.68		1.3											

SPECIES / SAMPLES	A1	A2	B1	B2	B3	B4	B5	C1	C2	C3	C4	C5	C6	C7	D1	D2	D3	D4	E3	E4	E5	E6	F1	F2	F3	F4	G1	G2	G3	G4	H	I	
depth (m)	0.43	1.30	3.21	2.43	2.40	1.41	0.90	4.38	4.78	4.46	3.40	2.68	1.73	n.o.	1.52	2.30	1.51	1.11	1.42	1.54	1.49	1.30	0.50	0.94	1.12	1.50	0.20	0.70	0.80	0.65	0.26	0.12	
<i>Neosidea mediterranea</i> (G.W. Müller, 1894)																		0.44	0.34														
<i>Neosidea</i> sp. 1										0.41						0.53																	
<i>Paracyclothera depressa</i> G.W. Müller, 1894		0.67				3.33				1.22		1.75			0.45		0.82			2.48	0.45		0.96									0.58	
<i>Paracyclothera</i> aff. <i>P. paralia</i> Babelo-Gonzalez, 1971										0.41																							
<i>Paracyclothera triquetra</i> (Reus, 1850)	4.00	0.67	0.59							1.63	1.23	1.17	2.11	1.84	0.90	1.59	3.27	7.38	1.31	1.35	0.35	2.24	2.37	1.92	0.85	1.95	4.94	2.63	1.21	1.85	3.99	1.16	
<i>Paracyclothera</i> aff. <i>P. flexuosa</i> (Brady, 1867)										0.41																							
<i>Paradoxostoma angustum</i> G.W. Müller, 1894										1.22							0.41	0.67									0.82						
<i>Paradoxostoma</i> aff. <i>P. angustum</i> G.W. Müller, 1894												1.05																					
<i>Paradoxostoma brevis</i> G.W. Müller, 1894										0.41																							
<i>Paradoxostoma caecum</i> G.W. Müller, 1894										0.81																							
<i>Paradoxostoma incongruens</i> G.W. Müller, 1894																		1.34															
<i>Paradoxostoma</i> aff. <i>P. porticum</i> Kie, 1942			1.76																														
<i>Paradoxostoma</i> aff. <i>P. rarum</i> G.W. Müller, 1894										0.41																							
<i>Paradoxostoma</i> aff. <i>P. rotundatum</i> G.W. Müller, 1894										0.41																							
<i>Paradoxostoma simile</i> G.W. Müller, 1894												1.17							0.44			0.45											
<i>Paradoxostoma</i> aff. <i>P. trise</i> G.W. Müller, 1894																			0.44					0.43									
<i>Paradoxostoma</i> sp. 1																					0.35												
<i>Paradoxostoma</i> sp. 1										0.41								0.67										0.41					
<i>Parasidea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)										0.41	0.41				0.45																		
<i>Portocypina spinosa</i> G.W. Müller, 1894										0.41								2.01															
<i>Portocypina acuminata</i> (G.W. Müller, 1894)										0.41					0.45														0.40				
<i>Portocypina turrida</i> (G.W. Müller, 1894)										0.41																							
<i>Procytheridea complicata</i> (Ruggieri, 1953)	4.70	0.67	4.55	2.65	10.00	2.86				0.41	1.23			3.07	1.81	1.06	0.41	0.67	0.44	0.34	0.35	0.45	1.18	0.96	1.28	1.30	1.65	4.45	0.81	2.31	9.78	18.02	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)			0.88											2.11	1.23		0.82	0.67		0.35			2.96		1.30	1.65	2.02	2.43	2.31	2.31	2.33		
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																							1.18										
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)	2.01										0.82			1.84	0.53							0.45		0.96	0.43						1.45		
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	
<i>Procytheridea</i> aff. <i>P. reticulata</i> (G.W. Müller, 1894)																																	




Fig. 5 - Ostracod abundance (A = number of specimens per gram of dry sediment) and diversity (SR = number of species, SI = Shannon index). Horizontal line indicates medium value.

abundance, *Leptocythere levis*, *L. affinis*, *L. rubritincta*, *C. adriatica*, *Pontocythere turbida*, *Neocytherideis mülleri*, *Urocythereis* sp. 1, *U. margaritifera*, *X. cypria*, *Callistocythere* aff. *C. gilva*, *Aurila punctata*, *Cytherelloidea sordida*.

The ostracod assemblages show a wide variability both in abundance and diversity (Fig. 5). Abundance ranges from 0.33 (sample C2) to 35.61 (sample H). The lowest diversity is recorded in the sample C1 (SR = 6, SI = 1.59), the highest in the sample C3 (SR = 75, SI = 3.65).

ASSEMBLAGE INTERPRETATION

Samples showing low values of abundance and diversity have been generally collected in the northern part of the bay, especially in coarse-grained sand (e.g. samples C1, C2, C6, B3, B4). In these sediments, assemblages are influenced by the relatively high-energy environment. In the southern, sheltered part of the bay, the bottom sediments are finer, and abundance and diversity values increase accordingly (Fig. 6). The most variable ecological index is abundance. Highest abundance occurs in the southernmost sample (H), which is the only collected in very fine sands.

Sample A1 is an exception to this trend, having very low abundance (A = 0.81) and low diversity (SR = 17, SI = 2.64) in fine sandy sediments. Location of this sample is about 50 m from the urbanized area, in a zone influenced by human pollution (Belmonte & Rubino, 1988). This factor seems to affect ostracod assemblages just for a few dozens of meters away the township. Indeed, the sample B1, at about 100 m from Porto Cesareo, shows rather high abundance (A = 16.96) and diversity (SR = 46, SI = 3.38).




Fig. 6 - Ostracod abundance, number of species and Shannon index in increasing order. Samples from the northern and southern part of the bay are, respectively, in grey and white colour.

Some well diversified assemblages have been recorded in transects C and D. These samples are located in the central-north sector of the bay, which is well connected with the open sea, with a depth of one meter at least.

The ostracod assemblages of the Porto Cesareo Lagoon comprise species which have been indicated by authors (*i.a.* G.W. Müller, 1894; Rome, 1964; Bonaduce et al., 1976; Athersuch, 1978; Yassini, 1979; Melis & Pugliese, 1985; Bonaduce et al., 1988; Athersuch et al., 1989) as phytophilous (Tab. 4) and typical of sandy bottom dwellers (Tab. 5).

Twenty-four species occur on any bottom type (Tab. 6). Few species show clear preferences for any substrate.

Phytophilous ostracod species	
<i>Aglaioocypris rara</i>	<i>Neonesidea longevaginata</i>
<i>Aurila convexa</i>	<i>Paracytheridea depressa</i>
<i>Callistocythere littoralis</i>	<i>Paradoxostoma</i> spp.
<i>Cytherelloidea sordida</i>	<i>Procytherideis complicata</i>
<i>Cytherois frequens</i>	<i>Propontocypris pirifera</i>
<i>Cytherois succinea</i>	<i>Sclerochilus gewemülleri</i>
<i>Loxoconcha rhomboidea</i>	<i>Semicytherura simplex</i>
<i>Neocytherideis mülleri</i>	<i>Xestoleberis cypria</i>
<i>Neonesidea corpulenta</i>	<i>Xestoleberis dispar</i>
<i>Neonesidea formosa</i>	<i>Xestoleberis plana</i>

Tab. 4 - Phytophilous ostracod species.

Ostracod species preferring sandy bottom	
<i>Carinocythereis whiteii</i>	<i>Loxoconcha rubritincta</i>
<i>Cytheretta adriatica</i>	<i>Semicytherura acuticostata</i>
<i>Cytheretta subradiosa</i>	<i>Semicytherura diafora</i>
<i>Heterocythereis voraginosa</i>	<i>Semicytherura sulcata</i>
<i>Leptocythere ramosa</i>	<i>Urocythereis margaritifera</i>

Tab. 5 - Ostracod species showing preference for sandy substrate.

Ostracod species occurring on any bottom type	
<i>Aurila convexa</i>	<i>Leptocythere levis</i>
<i>Aurila punctata</i>	<i>Loxoconcha stellifera</i>
<i>Basslerites berchoni</i>	<i>Neocytherideis mulleri</i>
<i>Callistocythere aff. C. gilva</i>	<i>Paracytheridea triquetra</i>
<i>Callistocythere intricatoides</i>	<i>Pontocythere turbida</i>
<i>Callistocythere sp. A</i>	<i>Semicytherura aff. S. inversa</i>
<i>Carinocythereis whiteii</i>	<i>Semicytherura sulcata</i>
<i>Cistacythereis (H.) turbida</i>	<i>Urocythereis distinguenda</i>
<i>Cyprideis torosa</i>	<i>Urocythereis sp. 1</i>
<i>Cytherelloidea sordida</i>	<i>Xestoleberis communis</i>
<i>Cytheretta adriatica</i>	<i>Xestoleberis cypria</i>
<i>Cytheretta subradiosa</i>	<i>Xestoleberis dispar</i>

Tab. 6 - Ostracod species recorded on any type of substrate.

Among the commonest species *A. punctata* and *A. convexa* dominate on coarser substrates. The absence of *L. gibberosa* and *U. margaritifera* in very fine sands is remarkable, as well as the rarity of *L. rubritincta* specimens on coarse-medium grained sands.

Data show that relationship between species distribution and type of bottom sediments is complex.

The ostracod assemblages are characterized by some species which are presently living exclusively or preferentially in the southern part of the Mediterranean

1 - <i>Xestoleberis cypria</i>	2 - <i>Loxoconcha pontica</i>	3 - <i>Microcythere cf. M. inflexa</i>	4 - <i>Aurila prasina</i>	5 - <i>Falsocythere terryi</i>	6 - <i>Callistocythere folliculosa</i>	7 - <i>Semicytherura trachina</i>	8 - <i>Urocythereis distinguenda</i>	9 - <i>Semicytherura aff. S. inversa</i>	10 - <i>"Eufrosina" minima</i>	Locality	References
+	+									Tripoli	Barra, 1997
+										Cyprus	Athersuch, 1979
	+	+	+	+						Tripoli	Bonaduce & Pugliese, 1975
	+	+								Naxos	Barbeito-Gonzalez, 1971
		+	+	+	+	+	+	+		Southern Adriatic Sea	Bonaduce <i>et al.</i> , 1976
			+							Gulf of Taranto	Bonaduce <i>et al.</i> , 1982
			+							Gulf of Gabès	Lachenal, 1989
				+	+					Egadi Islands	Melis & Pugliese, 1985
						+				Tunisian Shelf	Bonaduce <i>et al.</i> , 1988

Tab. 7 - References of "warm" ostracod species distribution reported in Fig. 7.

1 - <i>Semicytherura simplex</i>	2 - <i>Semicytherura ventricosa</i>	3 - <i>Paradoxostoma angustum</i>	Locality	References
		+	Malta Isle	Bonaduce & Masoli, 1970
		+	Saint George Bay	Bonaduce <i>et al.</i> , 1970
+	+		Naxos	Barbeito-Gonzalez, 1971
+			Egadi Islands	Melis & Pugliese, 1985
+	+	+	Gulf of Naples	G.W. Müller, 1894
		+	Southern Adriatic Sea	Bonaduce <i>et al.</i> , 1976

Tab. 8 - References of "warm" ostracod species distribution reported in Fig. 8.

Sea (Tabs. 7-9; Figs. 7-8). These records confirm the subtropical affinity of the Porto Cesareo Lagoon environment.




Fig. 7 - Distribution of "warm" ostracod species listed in Tab. 7, occurring exclusively in the central and southern Mediterranean, not recorded in the central Tyrrhenian sea.




Fig. 8 - Distribution of "warm" ostracod species listed in Tab. 8, occurring exclusively in the central and southern Mediterranean, including the central Tyrrhenian sea.

Few rare species, that is *M. (M.) fulva*, *E. curta*, *M. vitrea*, *M. (T.) angulosa*, and *S. acuminata* are generally reported as taxa typical of not very shallow (mainly lower infralittoral-circalittoral) waters (*i.a.* Bonaduce et al., 1976, 1977, 1979, 1988; Yassini, 1979). It can be inferred that species characteristic of higher water depth, representing a negligible fraction of the assemblages, are able to survive in a very shallow, sheltered marine environment. Our data suggest that the coexistence, in a fossil assemblage, of upper infralittoral and infra-circalittoral forms does not

necessarily indicate that shallower taxa have been displaced in a deeper environment. The occurrence of scattered specimens of these relatively "deep" species confirms that paleoecological investigations should be based on assemblage analysis, including state of preservation and the presence of different moults.

For 13 species we reported the first record in the Ionian Sea (Tab. 10).

A small number of species (*Callistocythere* aff. *C. gilva*, *Callistocythere* aff. *C. rastrifera*, *Callistocythere* sp. A, *Cytheroma* aff. *C. karadagensis*, *Semicytherura*

Costa bati	Heterocythereis voraginosa	Neonesidea corpulenta	Microcytherura (M.) fulva	Caudites calceolatus	Triebelina raripila	Neonesidea longevaginata	Neonesidea formosa	Procytherideis complicata	Tenedocythere prava	Locality	References
+	+	+	+	+	+	+	+	+	+	Southern Adriatic Sea	Bonaduce et al., 1976
+	+	+	+	+	+	+	+	+	+	Tripoli	Bonaduce & Pugliese, 1975
+	+	+	+	+	+	+	+	+	+	Tripoli	Barra, 1997
+	+	+	+	+	+	+	+	+	+	Tunisian Shelf	Bonaduce et al., 1988
+	+	+	+	+	+	+	+	+	+	Gulf of Taranto	Bonaduce et al., 1982
+	+	+	+	+	+	+	+	+	+	Egadi Islands	Melis & Pugliese, 1985
+	+	+	+	+	+	+	+	+	+	Gulf of Gabès	Bonaduce & Masoli, 1968
+	+	+	+	+	+	+	+	+	+	Northern Adriatic Sea	Bonaduce et al., 1976
+	+	+	+	+	+	+	+	+	+	Malta Isle	Bonaduce & Masoli, 1970
+	+	+	+	+	+	+	+	+	+	Saint George Bay	Bonaduce et al., 1970
+	+	+	+	+	+	+	+	+	+	Naxos	Barbeito-Gonzalez, 1971
+	+	+	+	+	+	+	+	+	+	Bay of Bou Ismail	Yassini, 1979
+	+	+	+	+	+	+	+	+	+	Cyprus	Athersuch, 1979
+	+	+	+	+	+	+	+	+	+	Gulf of Naples	G.W. Müller, 1894
+	+	+	+	+	+	+	+	+	+	Monaco	Rome, 1964
+	+	+	+	+	+	+	+	+	+	Tunisian Shelf	Bonaduce et al., 1979
+	+	+	+	+	+	+	+	+	+	Limski Kanal	Uffenorde, 1972
+	+	+	+	+	+	+	+	+	+	Northern Tyrrhenian	Bonaduce et al., 1977
+	+	+	+	+	+	+	+	+	+	Southern Spain	Aranky, 1987
+	+	+	+	+	+	+	+	+	+	Saline Gulf	Arbulla et al., 2001

Tab. 9 - Distribution data and references of ten ostracod species living preferentially (not exclusively) in the central and southern Mediterranean.

Species	Locality	References
<i>Cytherois</i> aff. <i>C. fischeri</i>	Limski Kanal	Uffenorde, 1972
<i>Cytherois joachinoi</i>	Sarno Plain (Pleist.)	Barra, 1992
<i>Cytherois pontica</i>	Black Sea	Marinov, 1966
<i>Cytheroma</i> cf. <i>C. karadagensis</i>	Naxos	Barbeito-Gonzalez, 1971
	Limski Kanal	Uffenorde, 1972
" <i>Elofsonia</i> " <i>minima</i>	Adriatic Sea	Bonaduce et al., 1976
<i>Microcythere</i> cf. <i>M. inflexa</i>	Naxos	Barbeito-Gonzalez, 1971
	Naxos	Barbeito-Gonzalez, 1971
<i>Loxoconcha pontica</i>	Tripoli	Bonaduce & Pugliese, 1975
		Barra, 1997
<i>Paracytheridea triquetra</i>	Limski Kanal	Uffenorde, 1972
	Adriatic Sea	Bonaduce et al., 1976
<i>Paradoxostoma</i> aff. <i>P. ponticum</i>	Naxos	Barbeito-Gonzalez, 1971
<i>Sclerochilus gewernülleri</i>	Gulf of Naples	G. W. Müller, 1894
	Monaco	Rome, 1964
	Aegean Sea	Puri et al., 1969
	Sea of Candia	Puri et al., 1969
	Naxos	Barbeito-Gonzalez, 1971
	Adriatic Sea	Bonaduce et al., 1976
	Egadi Islands	Melis & Pugliese, 1985
<i>Urocythereis distinguenda</i>	Adriatic Sea	Bonaduce et al., 1976
	Cyprus	Athersuch, 1979
<i>Urocythereis</i> sp. 1	Tripoli	Bonaduce & Pugliese, 1975
		Athersuch, 1977
		Barra, 1997
	Saline Gulf	Arbulla et al., 2001
<i>Xestoleberis cypria</i>	Cyprus	Athersuch, 1979
	Tripoli	Barra, 1997

Tab. 10 - List of ostracod species never recorded previously in the Ionian Sea.

aff. *S. inversa*, *Urocythereis* sp. 1) left in open nomenclature, are probably undescribed species and need further investigations.

CONCLUSIONS

The study of foraminiferal and ostracod assemblages evidenced distribution trends of the benthic microfaunas of the Porto Cesareo Lagoon.

Assemblages are characteristic of shallow warm water. The tropical/subtropical affinity of the inlet, previously assessed through the findings of algae, sabellid, echinoderm, gastropod, and sponge warm water species, is confirmed by the dominance of the foraminifer *Peneroplis pertusus*, the presence of *P. planatus*, *Spirolina acicularis* and *Glabratella patelliformis*. Numerous ostracod species show a geographical distribution restricted to the southern areas of the Mediterranean Sea.

Both foraminifers and ostracods are represented by species living on sandy substrate and phytophilous species. Epiphyte taxa prevail in the foraminifer assemblages.

Euryhaline foraminifers (*Ammonia beccarii*, *Haynesina depressula*) are relatively common in the southern part of the lagoon, probably due to karstic freshwater supply. Ostracod distribution in the lagoon seems not influenced by salinity variations, and testifies a fully marine environment.

The presence of *Ammonia gaimardi* and the low values of ostracod abundance and diversity in some samples located near the township of Porto Cesareo can indicate a certain influence of human pollution.

At the mouth of the lagoon, on sediments deposited in high energy waters, foraminifers are large, thick and frequently abraded; ostracod assemblages show low abundance and diversity.

The lack of abraded ostracod shells is probably due to their fragility. Actually, ostracod valves, compared with foraminifer tests, can be easily broken in high energy waters or dissolved within a short time in waters undersaturated in calcium carbonate. Some high-diversity ostracod assemblages have been recorded in the portion of the lagoon well connected to the sea, while the highest abundance was mainly recorded in the southern, sheltered sector.

Foraminifers have been divided in three groups: 1) species of open zone, 2) species of sheltered zone, and 3) ubiquitous species.

A little number of foraminifer and ostracod species, generally reported from deeper waters, occurs in the lagoon. Present data seem to demonstrate that these taxa are able to survive in the calm waters of the lagoon, even if the environmental parameters are probably not ideal for these species.

ACKNOWLEDGEMENTS

We especially acknowledge Piero De Castro who personally oversaw the sampling in the La Strea Bay. Our thanks also to Pasquale Raia who kindly reviewed the manuscript and the referees, Maria Gabriella Carboni (Università degli Studi di Roma

“La Sapienza”) and Nevio Pugliese (Università degli Studi di Trieste) for their appropriate remarks.

REFERENCES

- Aranki J.F. (1987). Marine Lower Pliocene Ostracoda of southern Spain with notes on the Recent fauna. *Bulletin of the Geological Institutions of the University of Uppsala - New Series*, 13: 1-94.
- Arbulla D., Pugliese N. & Russo A. (2001). Ostracodi del Golfo Saline (Sardegna nord-orientale). *Studi Trentini di Scienze Naturali - Acta Geologica*, 77: 25-35.
- Athersuch J. (1977). The genus *Urocythereis* (Crustacea: Ostracoda) in Europe, with particular reference to Recent Mediterranean species. *Bulletin of the British Museum of Natural History (Zoology)*, 32 (7): 247-283.
- Athersuch J. (1978). The genus *Xestoleberis* (Crustacea: Ostracoda) with particular reference to Recent Mediterranean species. *Pubblicazioni della Stazione Zoologica di Napoli*, 40: 282-343.
- Athersuch J. (1979). The ecology and distribution of the littoral ostracods of Cyprus. *Journal of Natural History*, 13: 135-160.
- Athersuch J., Horne D.J. & Whittaker J.E. (1989). Marine and Brackish Water Ostracods. In Kermack D.M. & Barnes R.S.K. (eds.), *Synopses of the British Fauna (new series)*, 43: 343 pp.
- Barbeito-Gonzalez P.J. (1971). Die Ostracoden des Küstenbereiches von Naxos (Griechenland) und ihre Lebensbereiche. *Mitteilungen aus dem hamburgischen zoologischen Museum und Institut*, 67: 255-326.
- Barra D. (1992). *Cytherois joachinoi*, nuova specie di ostracode marino del Pleistocene del Mediterraneo. *Atti del Museo Geologico e Paleontologico di Monfalcone*, 5: 3-11.
- Barra D. (1997). The shallow water marine ostracods of Tripoli (Lybia) and their geographical distribution in the Mediterranean. *Revista Española de Micropaleontología*, 29 (3): 71-106.
- Belmonte G. & Rubino G. (1988). Chimica e fisica delle acque. In Cinelli F., Cognetti G., Grasso M., Mongelli S., Orlando E. & Pagliai A.M. (eds.), *Studio ecologico dell'area marina di Porto Cesareo*, Congedo Editore: 31-47.
- Blanc-Vernet L. (1969). Contribution à l'étude des foraminifères de Méditerranée. Relations entre la microfaune et les sédiments; biocoenoses actuelles, thanatocoenoses Pliocènes et Quaternaires. *Recueil des travaux de la Station Marine d'Endoume, Marseille*, 64: 281 pp.
- Blanc-Vernet L. (1974). Microfaune de quelques dragages et carottages effectués devant les côtes de Tunisie (Golfe de Gabes) et de Libye. *Géologie Méditerranéenne*, 1: 9-26.
- Blanc-Vernet L. (1988). Repartition bathymétrique des foraminifères benthiques sur la plateforme continentale. *Géologie Méditerranéenne*, 15 (1): 5-12.
- Blanc-Vernet L., Clairefond P. & Orsolini P. (1979). Les foraminifères. In Burolet P.F., Clairefond P. & Winnock E. (eds.), *La mer Pélagienne. Géologie Méditerranée*, 6 (1): 171-209.
- Bonaduce G. & Masoli M. (1968). Marine Ostracoda of Tunisia. *Pubblicazioni della Stazione Zoologica di Napoli*, 36: 458-470.
- Bonaduce G. & Masoli M. (1970). Benthic marine Ostracoda from Malta. *Pubblicazioni della Stazione Zoologica di Napoli*, 38: 47-56.
- Bonaduce G., Brambati A. & Masoli M. (1970). Ostracoda and recent sediments of the Saint George Bay (Jun el Khudr, Lebanon). *Pubblicazioni della Stazione Zoologica di Napoli*, 38: 57-70.
- Bonaduce G., Ciampo G. & Masoli M. (1976). Distribution of Ostracoda in the Adriatic Sea. *Pubblicazioni della Stazione Zoologica di Napoli*, 40, supplemento 1: 1-304.
- Bonaduce G., Mascellaro P., Masoli M. & Pugliese N. (1982). Ostracodi. In Belfiore A., Bonaduce G., Garavelli C.,

- Mascellaro P., Masoli M., Mirabile L., Moncharmont M., Moretti M., Nuovo G., Pennetta M., Pescatore T., Placella P., Pugliese N., Russo B., Senatore M.R., Sgarrella F., Sansone E., Spezie G., Thorez J., Tramutoli M. & Vultaggio M., La sedimentazione recente del Golfo di Taranto (Alto Ionio, Italia). *Annali dell'Istituto Universitario Navale - Napoli*, 49-50, Appendice 3: 77-88.
- Bonaduce G., Masoli M., Minichelli G. & Pugliese N. (1979). The benthic ostracods. In Burolet P.F., Clairefond P. & Winnock E. (eds.), *La mer Pélagienne. Géologie Méditerranée*, 6 (1): 280-284.
- Bonaduce G., Masoli M. & Pugliese N. (1977). Ostracodi bentonici dell'alto Tirreno. *Studi Trentini di Scienze Naturali, Biologica*, 54: 243-261.
- Bonaduce G., Masoli M. & Pugliese N. (1988). Remarks on the Benthic Ostracoda on the Tunisian Shelf. In Hanai T., Ikeya N. & Ishizaki K. (eds.), *Evolutionary biology of Ostracoda its fundamentals and applications. Proceedings of the Ninth International Symposium on Ostracoda, held in Shizuoka, Japan 29 July-2 August 1985. Developments in Palaeontology and Stratigraphy*, 11: 449-466.
- Bonaduce G. & Pugliese N. (1975). Ostracoda from Lybia. *Pubblicazioni della Stazione Zoologica di Napoli*, 39: 129-135.
- Breman E. (1976). The distribution of ostracodes in the bottom sediments of the Adriatic Sea *Academisch Proefschrift*. 165 pp. Vrije Universiteit te Amsterdam, Amsterdam.
- Carrada G.C. & Fresi E. (1988). Le lagune salmastre costiere. Alcune riflessioni sui problemi e sui metodi. In Carrada G.C., Cicogna F. & Fresi E. (eds.), *Le lagune costiere: ricerca e gestione*, CLEM: 35-56.
- Castelli A., Curini Galletti M., Lardicci C., Giangrande A. & Variale Cognetti A.M. (1988). Comunità zoobentoniche. In Cinelli F., Cognetti G., Grasso M., Mongelli S. Orlando E. & Pagliai A.M. (eds.), *Studio ecologico dell'area marina di Porto Cesareo*, Congedo Editore: 72-86.
- Cita M.B. & Premoli Silva I. (1967). Sui foraminiferi incontrati in un pozzo perforato nella laguna di Venezia. *Memorie di Biogeografia Adriatica*, 7: 29-51.
- Colom G. (1971). Variabilité et stabilité des Peneroplinae (*Peneroplis*, *Dendritina*, *Spirolina*) depuis l'Eocène jusqu'aux temps actuels. *Revue de Micropaléontologie*, Numéro Spécial à la mémoire du Professeur Jean Cuvillier, 14 (5): 48-53.
- Coppa M.G. (1987). Les carottages holocènes de la plate-forme continentale sarde. *Géologie Méditerranéenne*, 14: 25-43.
- Coppa M.G. (1991). Un nuovo genere di foraminifero della piattaforma continentale sarda. In Bonfiglio, L. (ed.), *Celebrazioni del 1° centenario in memoria di Giuseppe Seguenza naturalista e paleontologo, Convegno di paleontologia e stratigrafia, Messina-Taormina 22-26 Maggio 1989. Atti della Accademia Peloritana dei Pericolanti, Classe di Scienze Fisiche Matematiche e Naturali*, 67 (1): 483-501.
- Coppa M.G. & Di Tuoro A. (1995). Preliminary data on the Holocene Foraminifera of the Cilento continental shelf (Tyrrhenian sea). *Revista Española de Paleontología*, 10 (2): 161-174.
- Coppa M.G., Russo B. & Siani G. (1994). The Holocene foraminiferal assemblages of the continental margin between Agropoli and Capo Palinuro (Tyrrhenian Sea, Italy). In Matteucci R., Carboni M.G. & Pignatti J.S. (eds.), *Studies on Ecology and Paleoecology of Benthic communities. Bollettino della Società Paleontologica Italiana, Volume Speciale*, 2: 67-91.
- Gherardi M., Giangrande A. & Corriero G. (2001). Epibiontic and endobiontic polychaetes of *Geodia cydonium* (*Porifera, Demospongiae*) from the Mediterranean Sea. *Hydrobiologia*, 443: 87-101.
- Lachenal A.M. (1989). Écologie des ostracodes du domaine méditerranéen: application au Golfe de Gabès (Tunisie orientale). Les variations du niveau marin depuis 30000 ans. *Documents des Laboratoires de Géologie de Lyon*, 108. 239 pp.
- Langer M. (1988). Recent epiphytic Foraminifera from Vulcano (Mediterranean Sea). In Benthos '86. Third International Symposium on Benthic Foraminifera, Genève, September 22-28, 1986. *Revue de Paléobiologie, Volume Spécial*, 2 (2): 827-832.
- Le Calvez J. & Le Calvez Y. (1958). Répartition des foraminifères dans la baie de Villefranche. 1-Miliolidae. *Annales de l'Institut Océanographique*, 35: 159-234.
- Loeblich A.R. Jr. & Tappan H. (1964). Sarcodina, chiefly "Thecamoebians" and Foraminifera. In Moore R.C. (ed.), *Treatise on Invertebrate Paleontology, Part C, Protista* 2. 900 pp. Geological Society of America and University of Kansas Press.
- Loeblich A.R. Jr. & Tappan H. (1987). Foraminiferal Genera and their Classification. 970 pp. Van Nostrand Reinhold Company, New York.
- Marinov T. (1966). *Cytherois pontica* n. sp. - eine neue Schwarzmeerostracode. *Crustaceana*, 10: 70-74.
- Mastroruzzi G., Palmentola G. & Ricchetti G. (1989). Aspetti della evoluzione olocenica della costa pugliese. *Memorie della Società Geologica Italiana*, 42: 287-300.
- Melis R. & Pugliese N. (1985). Le Ostracofaune bentoniche delle Isole Egadi (Sicilia). *Atti del Museo Geologico e Paleontologico di Monfalcone*, 3: 1-22.
- Mercurio M., Corriero G., Liaci L.S. & Gaino E. (2000). Silica content and spicule size variations in *Pellina semitubulosa* (*Porifera: Demospongiae*). *Marine Biology*, 137: 87-92.
- Müller G.W. (1894). Die Ostracoden des Golfes von Neapel und der angrenzenden Meeres-Abschnitte. *Fauna und Flora des Golfes von Neapel und der angrenzenden Meeres-Abschnitte, Herausgegeben von der Zoologischen Station zu Neapel*, 21 (1-8): 404 pp.
- Murray J.W. (1991). Ecology and Paleoecology of Benthic Foraminifera. 365 pp. Longman Scientific & Technical (John Wiley), Harlow.
- Pardi G., Salghetti Drioli U. & Della Pietà F. (1988). Vegetazione bentonica. In Cinelli F., Cognetti G., Grasso M., Mongelli S., Orlando E. & Pagliai A.M. (eds.), *Studio ecologico dell'area marina di Porto Cesareo*, Congedo Editore: 67-71.
- Parenzan P. (1983). Puglia Marittima. Aspetti geologici e biologia marina (1-2). 688 pp. Congedo Editore, Galatina.
- Passeri L. (1974). Sedimentazione carbonatica attuale e diagenesi precoce nella laguna di Porto Cesareo (Penisola Salentina). Atti della riunione sul tema: Sedimentologia delle rocce carbonatiche di mare sottile - Perugia 5, 6 e 7 ottobre 1973. *Bollettino della Società Geologica Italiana*, 92 (Supplemento): 3-40.
- Peres J.M. & Picard J. (1964). Nouveau manuel de bionomie benthique de la Mer Méditerranée. *Recueil des travaux de la Station Marine d'Endoume, Marseille*, Bull. 31 (47): 137 pp.
- Phleger F.B. (1960). Ecology and distribution of recent foraminifera. 297 pp., The Johns Hopkins Press, Baltimore.
- Puri H.S., Bonaduce G. & Gervasio A.M. (1969). Distribution of Ostracoda in the Mediterranean. In Neale J.W. (ed.), *The Taxonomy, Morphology and Ecology of Recent Ostracoda*, Oliver & Boyd: 358-411.
- Ricchetti G. & Nuovo G. (1988). Aspetti geologici e geomorfologici. In Cinelli F., Cognetti G., Grasso M., Mongelli S., Orlando E. & Pagliai A.M. (eds.), *Studio ecologico dell'area marina di Porto Cesareo*, Congedo Editore: 11-30.
- Rome D.R. (1964). Ostracodes des environs de Monaco, leur distribution en profondeur, nature des fonds marins explorés. In Ostracods as ecological and palaeoecological indicators. Simposio internazionale sotto gli auspici della Fondazione Antonio e Rinaldo Dohrn presso la Stazione Zoologica di Napoli, 10-19 Giugno 1963. *Pubblicazioni della Stazione Zoologica di Napoli*, 33 (Supplemento): 200-212.
- Scorziello R. (1990). Foraminiferi della spiaggia di Tripoli (Libia). In Robba E. (ed.), *Proceedings of the Fourth Symposium on Ecology and Paleoecology of Benthic Communities, Sorrento*

- 1-5 Novembre 1988, Museo Regionale di Scienze Naturali - Torino: 635-641.
- Seiglie G.A. & Bermudez P.J. (1965). Monografia de la familia de Foraminiferos Glabratellidae. *Geos*, 12: 15-64.
- Sgarrella F. & Barra D. (1985). Distribuzione dei Foraminiferi bentonici nel Golfo di Salerno (Basso Tirreno, Italia). *Bollettino della Società dei Naturalisti in Napoli*, 93: 1-58.
- Sgarrella F., Barra D. & Improta A. (1985). The benthic foraminifers of the Gulf of Policastro (Southern Tyrrhenian Sea, Italy). *Bollettino della Società dei Naturalisti in Napoli*, 92: 67-114.
- Sgarrella F. & Moncharmont Zei M. (1993). Benthic Foraminifera of the Gulf of Naples (Italy): systematics and autoecology. *Bollettino della Società Paleontologica Italiana*, 32 (2): 145-264.
- Uffenorde H. (1972). Oekologie und jahreszeitliche Verteilung rezenter bentonischer Ostracoden des Limski kanal bei Rovinj (nördliche Adria). *Göttinger Arbeiten zur Geologie und Paläontologie*, 13: 1-121.
- Vénec-Peyré M.T. & Le Calvez Y. (1988). Les Foraminifères épiphytes de l'herbier de posidonies de Banyuls-sur-Mer (Méditerranée occidentale): étude des variations spatiotemporelles du peuplement. *Cahiers de Micropaléontologie, nouvelle série*, 3 (2): 21-40.
- Yassini I. (1979). The littoral system Ostracodes from the bay of Bou-Ismaïl, Algiers, Algeria. *Revista Española de Micropaleontología*, 11 (3): 353-416.

Manuscript received 02 December 2005

Revised manuscript accepted 17 May 2006